

## DAEs - Control and Numerics

### Exercise Sheet 1 - Solution theory

**Exercise 1 (Regularity)**

Check whether the matrix pairs

$$\left( \begin{pmatrix} 1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix} \right), \quad \left( \begin{pmatrix} 2 & -1 & 1 \\ 3 & -2 & 2 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 1 & -1 & 1 \end{pmatrix} \right)$$

are regular or singular and determine their Kronecker canonical forms by elementary row and column transformations.

**Exercise 2 (Wong-sequences)**

Consider

$$A := \begin{pmatrix} 3 & 0 & 1 & 0 \\ 0 & 2 & 2 & -1 \\ 1 & 2 & 3 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}, \quad E := \begin{pmatrix} 1 & -1 & -3 & 0 \\ 0 & 2 & 0 & -1 \\ -3 & -1 & 1 & 2 \\ -2 & -2 & 0 & 2 \end{pmatrix}.$$

Calculate the so-called Wong sequences, given by

$$\begin{aligned} \mathcal{V}_0 &:= \mathbb{R}^n, & \mathcal{V}_{i+1} &:= A^{-1}(E\mathcal{V}_i), \\ \mathcal{W}_0 &:= \{0\}, & \mathcal{W}_{i+1} &:= E^{-1}(A\mathcal{W}_i), \end{aligned}$$

where  $M^{-1}(\mathcal{S}) := \{ x \in \mathbb{R}^n \mid Mx \in \mathcal{S} \}$  is the pre-image of the some set  $\mathcal{S} \subseteq \mathbb{R}^n$  under the matrix  $M \in \mathbb{R}^{n \times n}$  and  $M\mathcal{S} := \{ Mx \in \mathbb{R}^n \mid x \in \mathcal{S} \}$  is the image of  $\mathcal{S}$  under  $M$ .

Choose full rank matrices  $V, W$  with  $\text{im } V = \bigcap_{i \in \mathbb{N}} \mathcal{V}_i$  and  $\text{im } W = \bigcup_{i \in \mathbb{N}} \mathcal{W}_i$  and calculate  $PEQ$  and  $PAQ$  where  $P := [V, W]$  and  $Q := [EV, AW]^{-1}$ . What do you observe?

**Exercise 3 (Drazin inverse)**

Consider a regular matrix pair  $(E, A)$  and any  $\lambda \in \mathbb{R}$  such that  $\det(\lambda E - A) \neq 0$ . Prove that  $\hat{E} := (\lambda E - A)^{-1}E$  and  $\hat{A} := (\lambda E - A)^{-1}A$  commute, i.e.  $\hat{E}\hat{A} := \hat{A}\hat{E}$ .

Furthermore, calculate the Drazin inverse  $\hat{E}^D$  of  $\hat{E}$  and  $\hat{A}^D$  of  $\hat{A}$  with  $(E, A)$  as in Exercise 2 and  $\lambda = -1$ .

**Exercise 4 (A bad distribution)**

Denote with  $\mathcal{C}_0^\infty$  the space of smooth functions  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  with bounded support. Consider the following linear mapping

$$\mathcal{C}_0^\infty \ni \varphi \mapsto \sum_{i \in \mathbb{N}} \frac{(-1)^i}{i+1} \varphi\left(\frac{(-1)^i}{i+1}\right).$$

Show that this mapping is well defined for all  $\varphi \in \mathcal{C}_0^\infty$ , i.e. the infinite sum converges in  $\mathbb{R}$ .

Consider now the “restriction” of this mapping to the open interval  $(0, \infty)$ :

$$\mathcal{C}_0^\infty \ni \varphi \mapsto \sum_{i \in \mathbb{N}} \frac{1}{2i+1} \varphi\left(\frac{1}{2i+1}\right).$$

Show that this mapping is not well defined if  $\varphi(0) \neq 0$ .