

ON STABILITY OF LINEAR TIME-VARYING SECOND-ORDER DIFFERENTIAL EQUATIONS

BY

LUU HOANG DUC (*Fachbereich Mathematik, J.W. Goethe Universität, Frankfurt, Germany*),
ACHIM ILCHMANN (*Institut für Mathematik, Technische Universität Ilmenau, Ilmenau, Germany*),
STEFAN SIEGMUND (*Fachbereich Mathematik, J.W. Goethe Universität, Frankfurt, Germany*),

AND

PETER TARABA (*Fachbereich Mathematik, J.W. Goethe Universität, Frankfurt, Germany*)

Abstract. We derive sufficient conditions for stability and asymptotic stability of second order, scalar differential equations with differentiable coefficients.

1. Introduction. We study, for differentiable $a_0, a_1 : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$, stability properties of linear time-varying second-order differential equations of the form

$$\ddot{x} + a_1(t)\dot{x} + a_0(t)x = 0, \tag{1.1}$$

which describe, if a_0, a_1 are non-negative, an oscillator with damping a_1 and rigidity a_0 .

Numerous sufficient conditions for uniform (and asymptotic) stability of (1.1) are derived in terms of bounds of the coefficients and its derivatives. These results are presented in Section 2. In the remainder of the present section, we first recall and make precise

Received June 22, 2005.

2000 *Mathematics Subject Classification.* Primary 34A30, 34D20, 35B40 .

.
. .
. .

Current address: Fachbereich Mathematik, J.W. Goethe Universität, Robert-Mayer-Straße 10, 60325 Frankfurt, Germany.

E-mail address: `lhduc@math.uni-frankfurt.de`

Current address: Institut für Mathematik, Technische Universität Ilmenau, Weimarer Straße 25, 98693 Ilmenau, Germany.

E-mail address: `achim.ilchmann@tu-ilmenau.de`

Current address: Fachbereich Mathematik, J.W. Goethe Universität, Robert-Mayer-Straße 10, 60325 Frankfurt, Germany.

E-mail address: `siegmund@math.uni-frankfurt.de`

Current address: Fachbereich Mathematik, J.W. Goethe Universität, Robert-Mayer-Straße 10, 60325 Frankfurt, Germany.

E-mail address: `taraba@math.uni-frankfurt.de`

©XXXX Brown University

various concepts of stability, and then give a brief overview about the results available in the literature and related to our findings.

Writing $y = (y_1, y_2)^T = (x, \dot{x})^T$, equation (1.1) is equivalent to the planar first-order system

$$\dot{y} = A(t)y, \quad \text{where} \quad A(t) = \begin{bmatrix} 0 & 1 \\ -a_0(t) & -a_1(t) \end{bmatrix}. \quad (1.2)$$

For given $(x^0, x^1)^T = y^0 \in \mathbb{R}^2$ and $t_0 \geq 0$, we denote by $x(\cdot; t_0, x^0, x^1) : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$, $y(\cdot; t_0, y^0) : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^2$ the solution of (1.1), (1.2) with initial data

$$(x(t_0; t_0, x^0, x^1), \dot{x}(t_0; t_0, x^0, x^1)) = (x^0, x^1), \quad y(t_0; t_0, y^0) = y^0,$$

respectively. Uniqueness and existence of the solution on $\mathbb{R}_{\geq 0}$ is well known.

To investigate stability properties of the zero solution of (1.1), respectively of (1.2), or sufficient conditions of it, we recall the following definitions:

DEFINITION 1.1. The zero solution of (1.2), and equally (1.1), is said to be

- *stable* if, and only if, for every $t_0 \geq 0$ and $\varepsilon > 0$ there exists a $\delta = \delta(\varepsilon, t_0) > 0$ such that

$$\|y^0\| < \delta \implies \|y(t; t_0, y^0)\| < \varepsilon \quad \text{for all } t \geq t_0;$$

- *uniformly stable*, if, and only if, it is stable and the δ in (i) does not depend on t_0 ;
- *attractive* if, and only if, for all $(y^0, t_0) \in \mathbb{R}^2 \times \mathbb{R}_{\geq 0}$, we have $\lim_{t \rightarrow \infty} y(t; t_0, y^0) = 0$;
- *uniformly attractive* if, and only if, for all $\delta > 0$ there exists $T = T(\delta) > 0$ such that, for all $t_0 \geq 0$, for all $t \geq t_0 + T$, and for all $y^0 \in \mathbb{R}^2$ we have $\|y(t; t_0, y^0)\| \leq \delta \|y^0\|$;
- *(uniformly) asymptotically stable* if, and only if, it is (uniformly) stable and (uniformly) attractive, respectively.

REMARK 1.2.

- (a) Stability, (uniform) asymptotic stability and (uniform) attractivity can be defined for arbitrary solutions of system (1.2). By linearity of system (1.2) each solution has the same stability type and hence we may speak of stability properties of the system instead of stability properties of the zero solution.
- (b) We repeatedly use the fact that for linear systems attractivity implies stability, i.e. if system (1.2) is attractive then it is already asymptotically stable.
- (c) To prove stability, asymptotic stability or attractivity it suffices to check the conditions in Definition 1.1 for a single $t_0^* \geq 0$ instead of for all $t_0 \geq 0$, since one can use continuous dependence of the solution on the initial condition to estimate the size of $\|\lambda(t; t_0, y^0)\|$ on the compact interval $[\min\{t_0, t_0^*\}, \max\{t_0, t_0^*\}]$.

A first step in the direction of sufficient conditions for asymptotic stability is via the time depending eigenvalues of the matrix $A(t)$ in (1.2). However, in general these eigenvalues do not contain sufficient or necessary information about the stability type; Hoppenstead [7] gives a two dimensional example where both eigenvalues of $A(t)$ are located at -1 for all $t \geq 0$, however, the system has an unstable solution. Only if the eigenvalues of $A(t)$ depend “slowly” on time, then asymptotic stability is related to

the time-dependent eigenvalues; see for example Desoer [3], Rosenbrock [14], Solo [19], Rugh [15] and the references in the latter textbook.

A different approach using topological methods is the Sacker-Sell or dichotomy spectrum (see Siegmund [17]). The spectrum consists of so-called spectral intervals in \mathbb{R} which generalize the eigenvalue real parts to the nonautonomous situation. If the spectrum is negative system (1.2) is uniformly asymptotically stable. Since only in rare cases it is possible to compute the dichotomy spectrum directly from the coefficients a_0 and a_1 , we do not follow this approach but instead derive stability criteria based on the coefficients.

If the rigidity coefficient $a_0(\cdot) \equiv a_0 > 0$ is constant and positive, then many sufficient conditions for stability and attractivity of the zero solution of (1.1) are known.

One approach to derive asymptotic stability of (1.1) is via an appropriate Lyapunov-function and by invoking LaSalle's Invariance Principle. If the coefficients of (1.1) satisfy

$$a_0(\cdot) \equiv a_0 > 0 \quad \text{and} \quad a_1(t) \geq 0 \quad \forall t \geq 0, \quad (1.3)$$

then the derivative of $V(t) := \dot{x}(t)^2 + a_0 x(t)^2$ along the solution $x(\cdot) \equiv x(\cdot; t_0, x^0, x^1)$, $(x^0, x^1) \in \mathbb{R} \times \mathbb{R}$, yields, for all $t_0 \geq 0$,

$$\frac{d}{dt} V(t) = -2a_1(t) \dot{x}(t)^2 \leq 0, \quad \forall t \geq t_0,$$

whence the zero solution is uniformly stable. Moreover, if

$$\exists \underline{a}_1, \bar{a}_1 > 0 \quad \forall t \geq 0 : \quad 0 < \underline{a}_1 \leq a_1(t) \leq \bar{a}_1, \quad (1.4)$$

holds, then, for all $t \geq 0$, $\frac{d}{dt} V(t) \leq -2\underline{a}_1 \dot{x}(t)^2$ and so LaSalle's Invariance Principle yields that $\lim_{t \rightarrow \infty} \dot{x}(t) = 0$, and the ω -limit set of the solution is $\ddot{x} + a_0 x = 0$, and therefore $\lim_{t \rightarrow \infty} x(t) = \lim_{t \rightarrow \infty} \dot{x}(t) = 0$. This result is well known (see Theorem 1 in [10]) and was also stated in [1]:

THEOREM 1.3 (Levin and Nohel [10]). If the coefficients of (1.1) satisfy (1.3) and (1.4), then the zero solution of (1.1) is asymptotically stable.

The upper bound in condition (1.4) cannot be omitted as can be seen from the equation

$$\ddot{x} + (1 + a_0 + e^t)\dot{x} + a_0 x = 0 \quad (1.5)$$

with positive and constant $a_0 > 0$, since it admits the solution $t \mapsto x(t) = 1/a_0 + e^{-t}$, and therefore the zero solution is not attractive.

In the attempt to weaken condition (1.4) two cases have been distinguished.

$$\textit{small damping: } a_0(\cdot) \equiv a_0 > 0 \quad \text{and} \quad \exists \bar{a}_1 > 0 \quad \forall t \geq 0 : \quad 0 \leq a_1(t) \leq \bar{a}_1,$$

and

$$\textit{large damping: } a_0(\cdot) \equiv a_0 > 0 \quad \text{and} \quad \exists \underline{a}_1 > 0 \quad \forall t \geq 0 : \quad 0 < \underline{a}_1 \leq a_1(t).$$

In both cases, asymptotic stability is not guaranteed: for small damping there might be an oscillation so that the zero solution is not attractive; for large damping one has to exclude the phenomenon of overdamping, i.e. the existence of a solution x with $\lim_{t \rightarrow \infty} x(t) = x_* \neq 0$. An example of the latter is equation (1.5).

Smith [18] proved that the condition

$$A_1(t) := \int_0^t a_1(s) ds \rightarrow \infty \quad \text{as } t \rightarrow \infty$$

is necessary for the asymptotic stability of equation (1.1). For the case of *large damping* he even gave a necessary and sufficient condition.

THEOREM 1.4 (Smith [18]). Suppose the coefficients of the equation (1.1) satisfy large damping. Then the condition

$$\int_0^\infty e^{-A_1(t)} \int_0^t e^{A_1(s)} ds dt = \infty \quad (1.6)$$

is equivalent to asymptotic stability of the zero solution of equation (1.1).

Hatvani, Krisztin and Totik [6] proved that for any $c > 0$ condition (1.6) is equivalent to

$$\sum_{n=1}^\infty [A_1(nc)^{-1} - A_1((n-1)c)^{-1}]^2 = \infty \quad (1.7)$$

which is sometimes easier to check, as can be seen for the examples $a_1(t) = t$ which does, and $a_1(t) = t^2$ which does not satisfy condition (1.6). In general condition (1.7) is still difficult to verify. The following two theorems provide sufficient conditions in the cases of large and small damping.

THEOREM 1.5 (Artstein, Infante [1]). Suppose the coefficients of the equation (1.1) satisfy large damping and, in addition, for some constants $A, B > 0$,

$$\int_0^t a_1(s) ds \leq A + Bt^2 \quad \text{for all } t \geq 0.$$

Then the zero solution of (1.1) is attractive and therefore asymptotically stable. Moreover, the exponent 2 is optimal in the sense that it cannot be replaced by any $2 + \varepsilon$, $\varepsilon > 0$.

THEOREM 1.6 (Hatvani [5]). In the case of small damping,

$$\limsup_{t \rightarrow \infty} \left(t^{-2/3} \int_0^t a_1(s) ds \right) > 0$$

is sufficient for asymptotic stability of equation (1.1). Moreover, the exponent $-2/3$ is optimal in the sense that it cannot be replaced by any $\varepsilon - 2/3$, $\varepsilon > 0$.

There are many results on asymptotic stability of certain nonlinear equations. We merely mention a series of papers by Pucci and Serrin [11, 12, 13], see also the references therein. A typical result restricted to the linear case (1.1) is the following theorem.

THEOREM 1.7 (Pucci, Serrin [11]). Suppose there exist continuous functions $\sigma, \delta : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ so that $\sigma\delta$ is bounded but

$$\int_0^t \sigma(\tau) d\tau \rightarrow \infty \quad \text{as } t \rightarrow \infty,$$

and either σ is of bounded variation or $\log \sigma$ is uniformly Lipschitz on $\mathbb{R}_{\geq 0}$.
 If the coefficients of (1.1) satisfy

$$a_0(\cdot) \equiv a_0 > 0 \quad \text{and} \quad 0 \leq \sigma(t) \leq a_1(t) \leq \delta(t) \quad \forall t \geq 0,$$

then the zero solution of (1.1) is asymptotically stable.

Compared to the amount of literature on small and large damping, there are only a few results on the stability of (1.1) available where both coefficients $a_0(\cdot)$ and $a_1(\cdot)$ are allowed to be non-constant, see e.g. Ignatyev [9] and for the case of complex-valued coefficient functions Hovhannisyan [8] and the references therein.

THEOREM 1.8 (Ignatyev [9]). Suppose that the coefficients of (1.1) satisfy the following

$$\begin{aligned} \exists M > 0 \forall t \geq 0 : \quad & |\dot{a}_0(t)| + |a_1(t)| \leq M \\ \exists \underline{a} > 0 \forall t \geq 0 : \quad & 0 < \underline{a} \leq \dot{a}_0(t) + 2a_0(t)a_1(t). \end{aligned}$$

Then the zero solution of (1.1) is uniformly asymptotically stable.

2. Criteria for asymptotic stability. In this section we extend some of the results mentioned in the introduction. The statement of the first theorem is in the spirit of Theorem 1.8, but the uniform bound on the absolute value of \dot{a}_0 is not needed. However, the method of proof is different and we only show uniform stability and attractivity, not uniform asymptotic stability.

THEOREM 2.1. Suppose that the coefficients of (1.1) satisfy the following

$$\exists \underline{a} > 0 \forall t \geq 0 : \quad 0 < \underline{a}_0 \leq a_0(t) \leq \bar{a}_0, \quad (2.1)$$

$$\exists \bar{a}_1 > 0 \forall t \geq 0 : \quad |a_1(t)| \leq \bar{a}_1, \quad (2.2)$$

$$\exists \underline{a} > 0 \forall t \geq 0 : \quad 0 < \underline{a} \leq \dot{a}_0(t) + 2a_0(t)a_1(t). \quad (2.3)$$

Then the zero solution of (1.1) is asymptotically stable. In fact, it is attractive and uniformly stable.

REMARK 2.2. If the rigidity coefficient $a_0(\cdot) \equiv a_0$ is a positive constant, then conditions (2.1)–(2.3) specialize to

$$0 < \frac{\underline{a}}{2a_0} \leq a_1(t) \leq \bar{a}_1 \quad \forall t \geq 0,$$

and therefore Theorem 2.1 generalizes Theorem 1.3.

Proof of Theorem 2.1: Let $(x^0, x^1) \in \mathbb{R} \times \mathbb{R}$, and $x(\cdot) \equiv x(\cdot; 0, x^0, x^1)$ be the solution to (1.1) starting at time 0 in (x^0, x^1) . In view of (2.1) and (2.3), the derivative of

$$V(t) := \frac{1}{2} x(t)^2 + \frac{1}{2a_0(t)} \dot{x}(t)^2, \quad t \geq 0,$$

along the solutions of (1.1) satisfies, by omitting the argument t ,

$$\begin{aligned} \dot{V}(t) &= x\dot{x} + \frac{1}{a_0} \dot{x}\ddot{x} - \frac{\dot{a}_0}{2a_0^2} \dot{x}^2 = x\dot{x} + \frac{1}{a_0} \dot{x}(-a_1\dot{x} - a_0x) - \frac{\dot{a}_0}{2a_0^2} \dot{x}^2 \\ &= -\frac{1}{2a_0^2} (\dot{a}_0 + 2a_0a_1) \dot{x}^2 \leq -\frac{\underline{a}}{2\bar{a}_0^2} \dot{x}^2 \leq 0, \quad \forall t \geq 0, \end{aligned}$$

and so, by integration and invoking (2.1),

$$\frac{1}{2}x(t)^2 + \frac{1}{2\bar{a}_0}\dot{x}(t)^2 \leq V(t) \leq V(0) \leq \frac{1}{2}(x^0)^2 + \frac{1}{2\underline{a}_0}(x^1)^2 \quad \forall t \geq 0.$$

Therefore, the zero solution of (1.1) is uniformly stable.

Note also that, in view of (2.1) and (2.2), $\ddot{x} = -a_1\dot{x} - a_0x$ is bounded. Furthermore,

$$V(0) \geq V(0) - V(t) = -\int_0^t \dot{V}(\tau) d\tau \geq \frac{\underline{a}}{2\bar{a}_0^2} \int_0^t \dot{x}(\tau)^2 d\tau \quad \forall t \geq 0,$$

and so \dot{x} is square integrable. Finally, since \dot{x} is square integrability and uniform continuous, the latter is a consequence of boundedness of \ddot{x} , and we may apply Barbălat's lemma [2] which then yields $\lim_{t \rightarrow \infty} \dot{x}(t) = 0$.

Next we prove that $V(t) \rightarrow 0$ as $t \rightarrow \infty$, which then implies that $x(t) \rightarrow 0$ as $t \rightarrow \infty$. Supposing that this is not true, then due to the fact that $t \mapsto V(t)$ is monotonically non-increasing, there exists $\delta > 0$ such that $V(t) \geq \delta^2$ for all $t \geq 0$. Since $\dot{x}(t) \rightarrow 0$ as $t \rightarrow \infty$, there exists $t_1 \geq 0$ such that

$$x(t)^2 = 2V(t) - \frac{1}{a_0(t)}\dot{x}(t)^2 \geq 2V(t) - \frac{1}{\underline{a}_0}\dot{x}(t)^2 \geq \delta^2 \quad \forall t \geq t_1.$$

By continuity of x , this means that either $x(t) \geq \delta$ or $x(t) \leq -\delta$ for all $t \geq t_1$. Suppose that $x(t) \geq \delta$ for all $t \geq t_1$.

Then $\ddot{x}(t) = -a_1(t)\dot{x}(t) - a_0(t)x(t) \leq |\bar{a}_1||\dot{x}(t)| - \underline{a}_0\delta$ together with $\dot{x}(t) \rightarrow 0$ as $t \rightarrow \infty$ yields existence of $t_2 \geq t_1$ such that

$$\ddot{x}(t) \leq -\underline{a}_0\frac{\delta}{2} \quad \forall t \geq t_2.$$

Integration gives

$$\dot{x}(t) - \dot{x}(t_2) = \int_{t_2}^t \ddot{x}(\tau) d\tau \leq -\underline{a}_0\frac{\delta}{2}(t - t_2) \quad \forall t \geq t_2,$$

and, by repeated integration, we arrive at

$$\begin{aligned} \delta - x(t_2) \leq x(t) - x(t_2) &= \int_{t_2}^t \dot{x}(\tau) d\tau \leq \int_{t_2}^t [\dot{x}(t_2) - \underline{a}_0\frac{\delta}{2}(\tau - t_2)] d\tau \\ &= \dot{x}(t_2)(t - t_2) - \frac{\underline{a}_0\delta}{4}(t - t_2)^2 \quad \forall t \geq t_2. \end{aligned}$$

Since the right hand side of this inequality tends to $-\infty$ as $t \rightarrow \infty$ and the left hand side is bounded, this is a contradiction. The case $x(t) \leq -\delta$ for all $t \geq t_1$ yields a contradiction in a similar manner. Thus $V(t) \rightarrow 0$ as $t \rightarrow \infty$, proving that $x(t) \rightarrow 0$, i.e. attractivity, and the proof of the theorem is complete. \square

EXAMPLE 2.3. Consider the equation

$$\ddot{x} + \dot{x} + a_0(t)x = 0 \tag{2.4}$$

where a_0 denotes a differentiable approximation of \hat{a}_0 defined, for the sequence $t_n = \frac{n(n-1)}{2}$, $n \geq 2$, as follows:

$$\hat{a}_0(t) = \begin{cases} t + 1, & t \in [0, 1] \\ -\frac{n}{n^2-1} t + \frac{n^2}{2(n+1)} + 2, & t \in [t_n, t_n + n - \frac{1}{n}] \\ n t + 2 - \frac{n^2(n+1)}{2}, & t \in [t_n + n - \frac{1}{n}, t_{n+1}]. \end{cases}$$

See Figure 1. There exists certainly an approximation a_0 of \hat{a}_0 such that the assump-




FIG. 1. The essential derivative of \hat{a}_0 is unbounded.

tions (2.1)-(2.3) are satisfied and so by Theorem 2.1 the zero solution of (2.4) is attractive and uniformly stable. However, the approximation a_0 has unbounded derivative, and hence Theorem 1.8 cannot be applied.

The next theorem deals with time-dependent, sign indefinite but bounded $a_1(\cdot)$ (including the case of small damping, see Hatvani's Theorem 1.6), moreover, the coefficient $a_0(\cdot)$ is only assumed to be bounded but not necessarily bounded away from 0.

THEOREM 2.4. Suppose that the coefficients of (1.1) satisfy the following

$$\int_0^t a_0(\tau) d\tau \rightarrow \infty \quad \text{as } t \rightarrow \infty \quad (2.5)$$

$$\exists M, \bar{a}_0 > 0 \forall t \geq 0 : \quad 0 < a_0(t) \leq \bar{a}_0 \quad (2.6)$$

$$\exists M, \bar{a}_1 > 0 \forall t \geq 0 : \quad |a_1(t)| \leq \bar{a}_1, \quad \text{and} \quad |\dot{a}_1(t)| < M |a_0(t)| \quad (2.7)$$

$$\exists k > 0 \forall t \geq 0 : \quad 0 < k a_0(t)^2 \leq \dot{a}_0(t) + 2a_0(t)a_1(t). \quad (2.8)$$

Then the zero solution of (1.1) is asymptotically stable.

REMARK 2.5. If the rigidity coefficient $a_0(\cdot) = a_0$ is constant and positive, then the assumptions of Theorem 2.4 specialize to

$$\forall t \geq 0 : \quad \frac{k a_0}{2} \leq a_1(t) \leq \bar{a}_1 \quad \text{and} \quad a_0 |\dot{a}_1(t)| < M.$$

Proof of Theorem 2.4:

Set

$$\begin{aligned} b &:= \min \left\{ \frac{1}{2\bar{a}_1}, \frac{k}{M+4+4b^2\bar{a}_0+\bar{a}_1k} \right\}, \\ c(t) &:= \frac{2b^2a_0(t) - ba_1(t) + 1}{a_0(t)}, \quad \forall t \geq 0 \\ d(t) &:= -2b - \dot{c}(t) - 2ba_0(t)c(t) + 2a_1(t) + 1, \quad \forall t \geq 0. \end{aligned}$$

Note that

$$c(t) - b^2 \geq b^2 + \frac{1}{2a_0(t)} \quad \forall t \geq 0, \quad (2.9)$$

and since

$$b \leq \frac{k}{M+4+4b^2\bar{a}_0+\bar{a}_1k},$$

it follows that

$$(1 - b\bar{a}_1)k + b[-M - 4 - 4b^2\bar{a}_0] \geq 0,$$

and therefore

$$\begin{aligned} d(t) &= (1 - b\bar{a}_1(t)) \frac{\dot{a}_0(t) + 2a_0(t)a_1(t)}{a_0(t)^2} \\ &\quad + b \left(\frac{\dot{a}_1(t)}{a_0(t)} - 4 \right) - 6b^2a_1(t) - 4b^2a_0(t) \geq 0 \quad \forall t \geq 0. \end{aligned} \quad (2.10)$$

The derivative of

$$W(t) := (x(t) + b\dot{x}(t))^2 + 2(c(t) - b^2)\dot{x}(t)^2, \quad \forall t \geq 0,$$

along the solution of (1.1) yields

$$\dot{W}(t) = -2ba_0(t)W(t) - d(t)\dot{x}(t)^2 \leq -2ba_0(t)W(t) \leq 0 \quad \forall t \geq 0$$

and so, by integration,

$$W(t) \leq \exp \left(-2b \int_0^t a_0(\tau) d\tau \right) W(t_0) \quad \forall t \geq 0. \quad (2.11)$$

Equation (2.11) shows stability of the zero solution. Moreover, in view of (2.5), we arrive at $\lim_{t \rightarrow \infty} W(t) = 0$, and therefore, $x(t) + b\dot{x}(t) \rightarrow 0$, as well as $b^2\dot{x}^2(t) < (c - b^2)\dot{x}^2(t) \rightarrow 0$ as $t \rightarrow \infty$. Finally, this implies that $\dot{x}(t) \rightarrow 0$ and thus also $x(t) \rightarrow 0$ as $t \rightarrow \infty$, which proves attractivity of the zero solution of (1.1) and by Remark 1.2 (b) also asymptotic stability, and hence the proof of the theorem is complete. \square

EXAMPLE 2.6. It readily follows from Theorem 2.4 that the zero solution of the equation

$$\ddot{x} + \frac{1}{1+t}\dot{x} + \frac{1}{1+t}x = 0$$

is asymptotically stable.

In the following two theorems we use a time-dependent transformation to transform equation (1.1) into $\ddot{z} + b(t)z = 0$. We distinguish the cases $b(t) \geq 0$ and $b(t) \leq 0$, for all $t \geq 0$. In both cases this method allows us to treat unbounded coefficients.

THEOREM 2.7. Suppose that the coefficients of equation (1.1) satisfy

$$b(t) := a_0(t) - \frac{1}{2}\dot{a}_1(t) - \frac{1}{4}a_1(t)^2 > 0 \quad \forall t \geq 0, \quad (2.12)$$

$$\lim_{t \rightarrow \infty} a_1(t) e^{-\frac{1}{2} \int_0^t (a_1(\tau) - g(\tau)) d\tau} = 0, \quad (2.13)$$

$$\lim_{t \rightarrow \infty} \sqrt{b(t)} e^{-\frac{1}{2} \int_0^t (a_1(\tau) - g(\tau)) d\tau} = 0, \quad (2.14)$$

$$\int_0^t [a_1(\tau) - g(\tau)] d\tau \rightarrow \infty \quad \text{as } t \rightarrow \infty, \quad (2.15)$$

where, for all $t \geq 0$, $g(t) := \max\{0, -\dot{b}(t)/b(t)\}$. Then the zero solution of equation (1.1) is asymptotically stable.

REMARK 2.8. By Remark 1.2 (c) it suffices to check whether condition (2.12) holds only for all $t \geq t_0$ for some $t_0 \geq 0$.

Proof of Theorem 2.7: Let $(x^0, x^1) \in \mathbb{R} \times \mathbb{R}$ and $x(\cdot) \equiv x(\cdot; 0, x^0, x^1)$ be the solution to (1.1). The transformation

$$z(t) = x(t) e^{\frac{1}{2} \int_0^t a_1(\tau) d\tau}, \quad \forall t \geq 0,$$

yields that (1.1) is equivalent to

$$\ddot{z} + b(t)z = 0, \quad \forall t \geq 0 \quad (2.16)$$

with $z \not\equiv 0$. The derivative of

$$V(t) = \sqrt{z(t)^2 + \dot{z}(t)^2/b(t)}, \quad \forall t \geq 0$$

along the solution of (2.16) satisfies

$$\begin{aligned} 2\dot{V}(t)V(t) &= -\frac{\dot{b}(t)}{b(t)} \frac{\dot{z}(t)^2}{b(t)} \leq g(t) [V(t)^2 - z(t)^2] \\ &\leq g(t) V(t)^2, \quad \forall t \geq 0 \end{aligned} \quad (2.17)$$

and hence integration gives

$$V(t) \leq e^{\frac{1}{2} \int_0^t g(\tau) d\tau} V(0), \quad \forall t \geq 0,$$

whence

$$|x(t)| = |z(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \leq e^{\frac{1}{2} \int_0^t (g(\tau) - a_1(\tau)) d\tau} V(0), \quad \forall t \geq 0,$$

and so, invoking (2.15), we have $\lim_{t \rightarrow \infty} x(t) = 0$. Since

$$\begin{aligned} |\dot{x}(t)| &\leq |\dot{z}(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} + \frac{1}{2} |a_1(t)| |z(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \\ &\leq \left(\sqrt{b(t)} + \frac{1}{2} |a_1(t)| \right) V(0) e^{\frac{1}{2} \int_0^t (g(\tau) - a_1(\tau)) d\tau} \quad \forall t \geq 0, \end{aligned}$$

an application of (2.13), (2.14) gives $\lim_{t \rightarrow \infty} \dot{x}(t) = 0$. Together with Remark 1.2 (c), this proves attractivity of the zero solution of (1.1) and by Remark 1.2,(b) also asymptotic stability. This completes the proof of the theorem. \square .

COROLLARY 2.9. Suppose the coefficients of (1.1) satisfy $a_1(\cdot) \equiv a_1$ constant and $\dot{a}_0(t) \geq 0$ for all $t \geq 0$. Then $g(\cdot) \equiv 0$, and (2.13), (2.15) in Theorem 2.7 are trivially fulfilled and a sufficient condition for attractivity of the zero solution becomes

$$b(t) = a_0(t) - \frac{1}{4}a_1^2 > 0 \quad \forall t \geq 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} b(t)e^{-a_1 t} = 0.$$

Thus, we may choose, for $0 < \lambda < a_1$, a non-decreasing function $a_0(\cdot)$ with $\frac{a_1^2}{4} \leq a_0(t) \leq e^{\lambda t}$ for all $t \geq 0$. In this case, the zero solution of equation (1.1) is asymptotically stable.

EXAMPLE 2.10. Consider, for $\alpha \in (0, 1)$, the equation

$$\ddot{x} + 2(t+1)^{-\alpha} \dot{x} + x = 0. \quad (2.18)$$

Then a straightforward calculation yields the existence of some $t_0 \geq 0$ such that, $t \geq t_0$,

$$\begin{aligned} 1 + \alpha \geq b(t) &= 1 + \alpha(t+1)^{-\alpha-1} - (t+1)^{-2\alpha} > 0; \\ \dot{b}(t) &= -\alpha(\alpha+1)(t+1)^{-\alpha-2} + 2\alpha(t+1)^{-2\alpha-1} \\ &= \alpha(t+1)^{-\alpha-2}(2(t+1)^{1-\alpha} - \alpha - 1) > 0. \end{aligned}$$

In this case $g(t) = 0$ for $t \geq t_0$ and therefore, for any $t \geq t_0$,

$$\begin{aligned} \int_0^t (a_1(\tau) - g(\tau)) d\tau &= \int_0^{t_0} (a_1(\tau) - g(\tau)) d\tau + \int_{t_0}^t (a_1(\tau) - g(\tau)) d\tau \\ &= \int_0^{t_0} (a_1(\tau) - g(\tau)) d\tau + 2 \int_{t_0}^t (\tau+1)^{-\alpha} d\tau \\ &= \int_0^{t_0} (a_1(\tau) - g(\tau)) d\tau + \frac{2}{1-\alpha} ((t+1)^{1-\alpha} - (t_0+1)^{1-\alpha}). \end{aligned}$$

Hence, the suppositions of Theorem 2.7 are satisfied and the zero solution of equation (2.18) is asymptotically stable. Note that the criterion of Hatvani [5] cannot be applied to this equation if $\alpha > \frac{1}{3}$.

In the following theorem we treat the case $b(t) < 0$ for all $t \geq 0$.

THEOREM 2.11. Suppose that the coefficients in equation (1.1) satisfy

$$\beta(t) := \frac{1}{4}a_1(t)^2 + \frac{1}{2}\dot{a}_1(t) - a_0(t) > 0 \quad \forall t \geq 0, \quad (2.19)$$

$$\lim_{t \rightarrow \infty} a_1(t) \sqrt{\beta(t) + 1} e^{\int_0^t (h(\tau) - \frac{1}{2}a_1(\tau)) d\tau} = 0, \quad (2.20)$$

$$\lim_{t \rightarrow \infty} \sqrt{\beta(t) + 1} e^{\int_0^t (h(\tau) - \frac{1}{2}a_1(\tau)) d\tau} = 0, \quad (2.21)$$

where,

$$h(t) := \left| \frac{\dot{\beta}(t)}{4\beta(t)} \right| + \sqrt{\beta(t) + \left(\frac{\dot{\beta}(t)}{4\beta(t)} \right)^2} \quad \text{for all } t \geq 0.$$

Then the zero solution of (1.1) is asymptotically stable.

REMARK 2.12. By Remark 1.2 (c) it suffices to check whether condition (2.19) holds only for all $t \geq t_0$ for some $t_0 \geq 0$.

Proof of Theorem 2.11: Using the same notation as in the proof of Theorem 2.7, we see that $\beta = -b$ and so equation (1.1) is equivalent to $\ddot{z} - \beta(t)z = 0$ which is equivalent, under $y = (z, \dot{z})^T$ to its first order formulation

$$\dot{y} = \begin{bmatrix} 0 & 1 \\ \beta(t) & 0 \end{bmatrix} y.$$

Applying the transformation

$$y = T(t)u = \begin{bmatrix} 1 & 1 \\ \sqrt{\beta(t)} & -\sqrt{\beta(t)} \end{bmatrix} u$$

yields

$$\dot{u} = B(t)u, \quad \text{where } B = T^{-1}AT - T^{-1}\dot{T} = \begin{bmatrix} \sqrt{\beta} - \dot{\beta}/4\beta & \dot{\beta}/4\beta \\ \dot{\beta}/4\beta & -\sqrt{\beta} - \dot{\beta}/4\beta \end{bmatrix}.$$

Note that if $\lambda(t)$ denotes the eigenvalue of $B(t)$ with biggest absolute value, then $h(t) = |\lambda(t)|$ and, invoking that $B(t)$ is symmetric,

$$\frac{d}{dt} \|u(t)\| \leq \|\dot{u}(t)\| = \|B(t)u\| \leq h(t) \|u(t)\| \quad \text{for almost all } t \geq 0,$$

and so

$$\begin{aligned} \|y(t)\| &= \|T(t)u(t)\| \leq \|T(t)\| \|u(t)\| \\ &\leq \sqrt{2 + 2\beta(t)} \|u(0)\| e^{\int_0^t h(\tau) d\tau} \quad \text{for all } t \geq 0. \end{aligned}$$

Furthermore,

$$\begin{aligned} |x(t)| &\leq |z(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \\ &\leq \|y(t)\| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \\ &\leq \sqrt{2 - 2b(t)} \|u(0)\| e^{\int_0^t [h(\tau) - \frac{1}{2} a_1(\tau)] d\tau} \quad \text{for all } t \geq 0, \end{aligned}$$

and

$$\begin{aligned} |\dot{x}(t)| &\leq |\dot{z}(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} + \frac{1}{2} |a_1(t)| |z(t)| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \\ &\leq [1 + \frac{1}{2} |a_1(t)|] \|y(t)\| e^{-\frac{1}{2} \int_0^t a_1(\tau) d\tau} \\ &\leq [1 + \frac{1}{2} |a_1(t)|] \sqrt{2 + 2\beta(t)} \|u(0)\| e^{\int_0^t [h(\tau) - \frac{1}{2} a_1(\tau)] d\tau} \quad \text{for all } t \geq 0. \end{aligned}$$

The conclusion now follows from the assumptions. \square

EXAMPLE 2.13. The unbounded coefficients of the differential equation

$$\ddot{x} + t\dot{x} + (t-1)x = 0$$

satisfy the assumptions of Theorem 2.11, and therefore its zero solution is asymptotically stable.

The following two theorems are based on a more geometric point of view. In Theorem 2.14 we construct a polygon which is invariant under the dynamics forward in time and give conditions for stability of system (1.2) or equivalently equation (1.1). In Theorem 2.15 we strengthen the hypothesis and provide geometric conditions for asymptotic stability. For both theorems, we consider, for $r > 0$, the compact set S^r by the boundary given by the polygon

$$\begin{aligned} s_1^r &:= \{(y_1, y_2) \mid y_1 = r, y_2 \in [-r, 0]\} \\ s_2^r &:= \{(y_1, y_2) \mid y_1 \in [0, r], y_2 = -r\} \\ s_3^r &:= \{(y_1, y_2) \mid y_1 \in [-r, 0], y_2 = -r - y_1\}, \end{aligned}$$

the point symmetric lines

$$\hat{s}_i^r := \{(-y_1, -y_2) \mid (y_1, y_2) \in s_i^r\} \quad \text{for } i = 1, 2, 3.$$

and $0 \in \text{int } S^r$.




FIG. 2. The vector field points into the forward invariant polygon S^1 .

THEOREM 2.14. Suppose the coefficients of the equation (1.1) satisfy

$$\forall t \geq 0 : \quad 0 < a_0(t) \leq a_1(t) \tag{2.22}$$

$$\forall t \geq 0 : \quad 1 \leq a_1(t). \tag{2.23}$$

Then the zero solution of equation (1.1) is uniformly stable.

Proof of Theorem 2.14: First, we show that S^1 is forward invariant, i.e. for any initial data $(y_1^0, y_2^0) \in S^1$ it follows that the unique solution of (1.2) satisfies, for all $t \geq 0$, $(y_1(t), y_2(t)) \in S^1$. Invoking (2.22) and (2.23), the following implications show that the vector field given by the right hand side of (1.2) points into S^1 along the polygon $s_1 \cup s_2 \cup s_3$. We omit the superscript “r” for $r = 1$ and the argument t in the following (see also Figure 2).

$$\begin{aligned}
 (y_1, y_2) = (1, 0) &\implies \dot{y}_1 = 0, \dot{y}_2 = -a_0 < 0 \\
 (y_1, y_2) \in s_1 &\implies \dot{y}_1 = y_2 \leq 0, \dot{y}_2 = -a_0 - a_1 y_2 \in \mathbb{R} \\
 (y_1, y_2) = (1, -1) &\implies \dot{y}_1 = -1, \dot{y}_2 = -a_0 + a_1 \geq 0 \\
 (y_1, y_2) \in s_2 &\implies \dot{y}_1 = -1, \dot{y}_2 = -a_0 y_1 + a_1 \geq -a_0 + a_1 \geq 0 \\
 (y_1, y_2) = (0, -1) &\implies (\dot{y}_1, \dot{y}_2) = (-1, a_1) \text{ with } a_1 \geq 1 \\
 (y_1, y_2) = (-1, 0) &\implies (\dot{y}_1, \dot{y}_2) = (0, a_0) \text{ with } a_0 > 0 \\
 (y_1, y_2) \in s_3 &\implies \dot{y}_1 = -1 - y_1, \dot{y}_2 \geq a_0 - a_1[-1 - y_0]
 \end{aligned}$$

By point symmetry, the vector field of (1.2) is also pointing inwards along the polygon $\hat{s}_1 \cup \hat{s}_2 \cup \hat{s}_3$. Therefore, S^1 is forward invariant and since (1.2) is linear, it follows that the zero solution of (1.2) is uniformly stable. This completes the proof of the theorem. \square

THEOREM 2.15. Suppose there exist $\varepsilon, M > 0$ such that the coefficients of equation (1.1) satisfy

$$\forall t \geq 0 : \quad 0 < \varepsilon \leq a_0(t) \leq a_1(t) \leq M \quad (2.24)$$

$$\forall t \geq 0 : \quad 1 + \varepsilon \leq a_1(t). \quad (2.25)$$

Then the zero solution of equation (1.1) is asymptotically stable.

Proof of Theorem 2.15: By Theorem 2.14 and Remark 1.2 (b), it suffices to prove attractivity of the zero solution. We proceed in two steps.

(i) If $y(\cdot)$ crosses the y_1 -axis only finitely many times, then there exists $t_0 \geq 0$ such that, for all $t \geq t_0$,

$$(a) \quad (y_1(t), y_2(t)) \in S^r \cap (\mathbb{R} \times \mathbb{R}_{\geq 0})$$

or

$$(b) \quad (y_1(t), y_2(t)) \in S^r \cap (\mathbb{R} \times \mathbb{R}_{\leq 0}).$$

We prove that in case of (b) the claim of the theorem follows, case (a) is proved analogously and omitted for brevity.

Since $\dot{y}_1(t) = y_2(t) \leq 0$ for all $t \geq t_0$ and y_1 is bounded, there exists $\widetilde{M} > 0$ such that

$$0 \leq \int_{t_0}^t -y_2(\tau) d\tau = y_1(t_0) - y_1(t) \leq \widetilde{M} \quad \forall t \geq t_0.$$

Therefore, $y_2 \in L^1$, and applying boundedness of y_1 and y_2 together with (2.24) to

$$|\dot{y}_2(t)| = |a_0(t)y_1(t) + a_1(t)y_2(t)| \leq M(|y_1(t)| + |y_2(t)|)$$

gives $\dot{y}_2 \in L^\infty$. So we may apply Barbălat’s lemma (see [2]) to conclude that $y_2(t) \rightarrow 0$ as $t \rightarrow \infty$.




FIG. 3. Solutions in S^r starting in the 4th quadrant do not cross the line L .

By monotonicity of y_1 , there exists $y_1^* \in \mathbb{R}$ such that $y_1(t) \rightarrow y_1^*$ as $t \rightarrow \infty$. Applying (2.24) again gives

$$\exists t_1 \geq t_0 \forall t \geq t_1 : \frac{\varepsilon}{2} |y_1^*| = |\dot{y}_2| \leq 2M |y_1^*|.$$

Since $y_2(t) \rightarrow 0$ as $t \rightarrow \infty$, it follows that $y_1^* = 0$.

(ii) Since the vector field along $(0, r) \times \{0\}$ is $\dot{y}_2 = -a_1 y_1 < 0$, $\dot{y}_1 = 0$, and along $(-r, 0) \times \{0\}$ is $\dot{y}_2 = -a_1 y_1 > 0$, $\dot{y}_1 = 0$, it remains to consider the case that there exists a sequence $(t_n)_{n \in \mathbb{N}}$, with

$$t_n < t_{n+1} \quad \text{and} \quad y_2(t_n) = 0, \quad y_1(t_n) > 0 \quad \forall n \in \mathbb{N}.$$

In the proof of Theorem 2.14 we have shown that S^r , $r > 0$, is positively invariant. Observe that, by (2.24) and (2.25), the vector field along the segment

$$L := \{(y_1, y_2) \in (-r/(1+\varepsilon), 0) \times (0, -r) \mid y_2 = -(1+\varepsilon)y_1 - r\}$$

satisfies

$$(\dot{y}_1, \dot{y}_2) \in (-r, 0) \times \left(\frac{a_0}{1+\varepsilon}, a_1 \right)$$

and consequently a solution starting in the polygon

$$L \cup \left(\left[\frac{-r}{1+\varepsilon}, r \right] \times \{0\} \right) \cup (\{r\} \times [-r, 0]) \cup ([0, r] \times \{-r\})$$

can leave only through the segment $\left[\frac{-r}{1+\varepsilon}, 0 \right] \times \{0\}$ (see also Figure 3). We may choose $t_0 \geq 0$ such that $y_2(t_0) = 0$ and $r := y_1(t_0)$. By the above observations, we also may choose a sequence $(t_n)_{n \in \mathbb{N}}$ such that $y_2(t_n) = 0$ and

$$0 < y_1(t_{n+1}) \leq \frac{1}{1+\varepsilon} y_1(t_n).$$

Together with the fact that

$$|y_2(t)| \leq |y_1(t_n)| \quad \text{for } t_n \leq t \leq t_{n+1}$$

(see also Figure 3) this implies that $y(t) \rightarrow 0$ as $t \rightarrow \infty$, and the proof of the theorem is complete. \square

Condition (2.25) ensures that each time a solution completes a rotation around the origin the norm is reduced by a factor less than 1. Whereas this condition might not be necessary for asymptotic stability, condition (2.24) cannot be omitted as the following examples show.

EXAMPLE 2.16. The condition $0 < \varepsilon < a_0(t)$ for all $t \geq 0$ in Theorem 2.15 cannot be omitted as the equation

$$\ddot{x} + (2 + e^{-t})\dot{x} + e^{-t}x = 0$$

shows. It has the solution $t \mapsto x(t) = 1 + e^{-t}$ which does not converge to 0 for $t \rightarrow \infty$.

EXAMPLE 2.17. The condition $\|a_1\|_\infty \in \mathbb{R}$ in Theorem 2.15 cannot be omitted as the equation

$$\ddot{x} + (2 + e^t)\dot{x} + x = 0$$

shows (cp. equation (1.5) with $a_0 = 1$). It has the solution $t \mapsto x(t) = 1 + e^{-t}$ which does not converge to 0 for $t \rightarrow \infty$.

REFERENCES

- [1] Z. Artstein, E.F. Infante, On the asymptotic stability of oscillators with unbounded damping. *Quart. Appl. Math.* **34** (1976/77), no. 2, 195–199.
- [2] I. Barbălat, Systèmes d'équations différentielles d'oscillations nonlinéaires, *Revue de Mathématiques Pures et Appliquées, Bucharest IV* (1959), 267–270.
- [3] C.E. Desoer, Slowly varying system $\dot{x} = A(t)x$. *IEEE Trans. Automatic Control* **14** (1969), 780–781.
- [4] L. Dieci, E. Van Vleck, Lyapunov and other spectra: a survey. Collected lectures on the preservation of stability under discretization (Fort Collins, CO, 2001), 197–218, SIAM, Philadelphia, PA, 2002.
- [5] L. Hatvani, Integral conditions on the asymptotic stability for the damped linear oscillator with small damping. *Proc. Amer. Math. Soc.* **124** (1996), 415–422.
- [6] L. Hatvani, T. Krisztin, V. Totik, A necessary and sufficient condition for the asymptotic stability of the damped oscillator. *J. Differential Equations* **119** (1995), 209–223.
- [7] F.C. Hoppenstaed, Singular perturbations on the infinite interval. *Trans. Am. Math. Soc.* **123** (1966), 521–535.
- [8] G.R. Hovhannisyan, Asymptotic stability for second-order differential equations with complex coefficients. *Electron. J. Differential Equations* **85** (2004), 20 pp.
- [9] A.O. Ignatyev, Stability of a linear oscillator with variable parameters. *Electron. J. Differential Equations* **17** (1997), 6 pp.
- [10] J.J. Levin, J.A. Nohel, Global asymptotic stability for nonlinear systems of differential equations and applications to reactor dynamics. *Arch. Rational Mech. Anal.* **5** (1960), 194–211.

- [11] P. Pucci, J. Serrin, Precise damping conditions for global asymptotic stability for nonlinear second order systems. *Acta Math.* **170** (1993), no. 2, 275–307.
- [12] P. Pucci, J. Serrin, Precise damping conditions for global asymptotic stability for nonlinear second order systems. II. *J. Differential Equations* **113** (1994), no. 2, 505–534.
- [13] P. Pucci, J. Serrin, Asymptotic stability for intermittently controlled nonlinear oscillators. *SIAM J. Math. Anal.* **25** (1994), no. 3, 815–835.
- [14] H.H. Rosenbrock, The stability of linear time-dependent control systems. *J. Electronics Control* **15** (1963), 73–80.
- [15] Rugh W.J. (1996), *Linear System Theory* 2nd ed., Prentice-Hall, Upper Saddle River, New Jersey
- [16] R.J. Sacker, G.R. Sell, A Spectral Theory for Linear Differential Systems. *J. Differential Equations* **27** (1978), 320–358.
- [17] S. Siegmund, Dichotomy spectrum for nonautonomous differential equations. *J. Dynam. Differential Equations* **14** (2002), no. 1, 243–258.
- [18] R.A. Smith, Asymptotic stability of $x'' + a(t)x' + x = 0$. *Quart. J. Math. Oxford Ser. (2)* **12** (1961), 123–126.
- [19] V. Solo, On the stability of slowly time-varying linear systems. *Math. Control Signals Systems* **7** (1994), no. 4, 331–350.